

# UMBRAL CALCULUS AND SPECIAL POLYNOMIALS

TAEKYUN KIM<sup>1</sup> AND DAE SAN KIM<sup>2</sup>

ABSTRACT. In this paper, we consider several special polynomials related to associated sequences of polynomials. Finally, we give some new and interesting identities of those polynomials arising from transfer formula for the associated sequences.

## 1. INTRODUCTION

In this paper, we assume that  $\lambda \in \mathbb{C}$  with  $\lambda \neq 1$ . For  $\alpha \in \mathbb{R}$ , the Frobenius-Euler polynomials are defined by the generating function to be

$$\left(\frac{1-\lambda}{e^t-\lambda}\right)^\alpha e^{xt} = \sum_{n=0}^{\infty} H_n^{(\alpha)}(x|\lambda) \frac{t^n}{n!}, \quad (\text{see } [1,5,13,15,20,22,23]). \quad (1.1)$$

In the special case,  $x = 0$ ,  $H_n^{(\alpha)}(0|\lambda) = H_n^{(\alpha)}(\lambda)$  are called the  $n$ -th *Frobenius-Euler numbers* of order  $\alpha$ . As is well known, the Bernoulli polynomials of order  $\alpha$  are given by

$$\left(\frac{t}{e^t-1}\right)^\alpha e^{xt} = \sum_{n=0}^{\infty} B_n^{(\alpha)}(x) \frac{t^n}{n!}, \quad (\text{see } [2,3,4,6,14,15,19,21]). \quad (1.2)$$

In the special case,  $x = 0$ ,  $B_n^{(\alpha)}(0) = B_n^{(\alpha)}$  are called the  $n$ -th *Bernoulli numbers* of order  $\alpha$ .

For  $n \geq 0$ , the *Stirling numbers of the second kind* are defined by generating function to be

$$(e^t - 1)^n = n! \sum_{l=n}^{\infty} S_2(l, n) \frac{t^l}{l!}, \quad (\text{see } [8-12,17,18]), \quad (1.3)$$

and the *Stirling numbers of the first kind* are given by

$$(x)_n = x(x-1) \cdots (x-n+1) = \sum_{l=0}^n S_1(n, l) x^l, \quad (\text{see } [7,8,10,17,18]). \quad (1.4)$$

Let  $\mathcal{F}$  be the set of all formal power series in the variable  $t$  over  $\mathbb{C}$  with

$$\mathcal{F} = \left\{ f(t) = \sum_{k=0}^{\infty} \frac{a_k}{k!} t^k \mid a_k \in \mathbb{C} \right\}. \quad (1.5)$$

Let  $\mathbb{P}$  be the algebra of polynomials in the variable  $x$  over  $\mathbb{C}$  and  $\mathbb{P}^*$  be the vector space of all linear functionals on  $\mathbb{P}$ . As a notation, the action of the linear functional

---

1991 *Mathematics Subject Classification.* 05A10, 05A19.

*Key words and phrases.* Bernoulli polynomial, Euler polynomial, Abel polynomial.

$L$  on a polynomial  $p(x)$  is denoted by  $\langle L | p(x) \rangle$ . Let  $f(t) = \sum_{k=0}^{\infty} a_k \frac{t^k}{k!} \in \mathcal{F}$ . Then we define the linear functional  $f(t)$  on  $\mathbb{P}$  by

$$\langle f(t) | x^n \rangle = a_n, \quad (n \geq 0), \quad (\text{see [10,12,16,17,18]}). \quad (1.6)$$

From (1.6), we note that

$$\langle t^k | x^n \rangle = n! \delta_{n,k}, \quad (n, k \geq 0), \quad (1.7)$$

where  $\delta_{n,k}$  is the Kronecker symbol (see [8, 10, 11, 17, 18]).

Let  $f_L(t) = \sum_{k=0}^{\infty} \frac{\langle L | x^k \rangle}{k!} t^k$ . Then, by (1.7), we get  $\langle f_L(t) | x^n \rangle = \langle L | x^n \rangle$ . The map  $L \mapsto f_L(t)$  is a vector space isomorphism from  $\mathbb{P}^*$  onto  $\mathcal{F}$ . Henceforth,  $\mathcal{F}$  thought of as both a formal power series and a linear functional. We shall call  $\mathcal{F}$  the *umbral algebra*. The umbral calculus is the study of umbral algebra (see [10, 16, 17, 18]).

The order  $o(f(t))$  of the non-zero power series  $f(t)$  is the smallest integer  $k$  for which the coefficient of  $t^k$  does not vanish (see [10, 11, 12, 17, 18]). If  $o(f(t)) = 1$ , then  $f(t)$  is called a *delta series*, and if  $o(f(t)) = 0$ , then  $f(t)$  is called an *invertible series*. Let  $o(f(t)) = 1$  and  $o(g(t)) = 0$ . Then there exists a unique sequence  $S_n(x)$  of polynomials such that  $\langle g(t)f(t)^k | S_n(x) \rangle = n! \delta_{n,k}$  where  $n, k \geq 0$ . The sequence  $S_n(x)$  is called *Sheffer sequence* for  $(g(t), f(t))$ , which is denoted by  $S_n(x) \sim (g(t), f(t))$ . If  $S_n(x) \sim (1, f(t))$ , then  $S_n(x)$  is called the *associated sequence* for  $f(t)$  (see [10, 16, 17, 18]). From (1.7), we note that  $\langle e^{yt} | p(x) \rangle = p(y)$ .

Let  $f(t) \in \mathcal{F}$  and  $p(x) \in \mathbb{P}$ . Then we have

$$f(t) = \sum_{k=0}^{\infty} \frac{\langle f(t) | x^k \rangle}{k!} t^k, \quad p(x) = \sum_{k=0}^{\infty} \frac{\langle t^k | p(x) \rangle}{k!} x^k, \quad (\text{see [17,18]}). \quad (1.8)$$

From (1.9), we can derive the following equation:

$$p^{(k)}(0) = \langle t^k | p(x) \rangle \quad \text{and} \quad \langle 1 | p^{(k)}(x) \rangle = p^{(k)}(0), \quad (\text{see [10,16,17,18]}). \quad (1.9)$$

for  $k \geq 0$ , by (1.9), we easily see that  $t^k p(x) = p^{(k)}(x) = \frac{d^k p(x)}{dx^k}$ .

Let  $S_n(x) \sim (g(t), f(t))$ . Then we see that

$$\frac{1}{g(\bar{f}(t))} e^{y\bar{f}(t)} = \sum_{k=0}^{\infty} \frac{S_k(x)}{k!} t^k, \quad \text{for all } y \in \mathbb{C}, \quad (1.10)$$

where  $\bar{f}(t)$  is the compositional inverse of  $f(t)$  (see [17, 18]).

Let  $p_n(x) \sim (1, f(t))$ ,  $q_n(x) \sim (1, g(t))$ . Then, the transfer formula for the associated sequence is given by

$$q_n(x) = x \left( \frac{f(t)}{g(t)} \right)^n x^{-1} p_n(x), \quad (\text{see [11,12,16,17,18]}). \quad (1.11)$$

For  $n \geq 0$ ,  $b \neq 0$ , the *Abel sequences* are given by

$$A_n(x; b) = x(x - bn)^{n-1} \sim (1, te^{bt}). \quad (1.12)$$

In this paper, we consider several special polynomials related to associated sequences of polynomials. Finally, we give some new and interesting identities of those polynomials arising from transfer formula for the associated sequences.

## 2. UMBRAL CALCULUS AND SPECIAL POLYNOMIALS

From (1.1), we note that

$$H_n^{(\alpha)}(x|\lambda) \sim \left( \left( \frac{e^t - \lambda}{1 - \lambda} \right)^\alpha, t \right). \quad (2.1)$$

Thus, we get

$$H_n^{(\alpha)}(x|\lambda) = \left( \frac{1 - \lambda}{e^t - \lambda} \right)^\alpha x^n. \quad (2.2)$$

Let us assume that

$$p_n(x) \sim (1, t(e^t - \lambda)), \quad q_n(x) \sim \left( 1, \left( \frac{e^t - \lambda}{1 - \lambda} \right)^a t \right), \quad (a \neq 0). \quad (2.3)$$

From  $x^n \sim (1, t)$ , (1.11) and (2.3), we note that

$$\begin{aligned} p_n(x) &= x \left( \frac{t}{t(e^t - \lambda)} \right)^n x^{-1} x^n = \frac{x}{(1 - \lambda)^n} \left( \frac{1 - \lambda}{e^t - \lambda} \right)^n x^{n-1} \\ &= \frac{1}{(1 - \lambda)^n} x H_{n-1}^{(n)}(x|\lambda). \end{aligned} \quad (2.4)$$

and

$$q_n(x) = x \left( \frac{1 - \lambda}{e^t - \lambda} \right)^{na} x^{-1} x^n = x H_{n-1}^{(an)}(x|\lambda). \quad (2.5)$$

From (1.11), (2.3), (2.4) and (2.5), we can derive

$$\begin{aligned} &\frac{1}{(1 - \lambda)^n} x H_{n-1}^{(n)}(x|\lambda) \\ &= x \left( \frac{t \left( \frac{e^t - \lambda}{1 - \lambda} \right)^a}{t(e^t - \lambda)} \right)^n x^{-1} x H_{n-1}^{(an)}(x|\lambda) \\ &= \frac{x}{(1 - \lambda)^{an}} (e^t - \lambda)^{(a-1)n} H_{n-1}^{(an)}(x|\lambda) \\ &= \frac{x}{(1 - \lambda)^{an}} \sum_{l=0}^{(a-1)n} \binom{(a-1)n}{l} (-\lambda)^{(a-1)n-l} e^{lt} H_{n-1}^{(an)}(x|\lambda) \\ &= \frac{x}{(1 - \lambda)^{an}} \sum_{l=0}^{(a-1)n} \binom{(a-1)n}{l} (-\lambda)^{(a-1)n-l} H_{n-1}^{(an)}(x + l|\lambda), \end{aligned} \quad (2.6)$$

where  $a, n \in \mathbb{N}$ . Therefore, by (2.6), we obtain the following theorem.

**Theorem 2.1.** *For  $a, n \in \mathbb{N}$ , we have*

$$H_{n-1}^{(n)}(x|\lambda) = \frac{1}{(1 - \lambda)^{(a-1)n}} \sum_{l=0}^{(a-1)n} \binom{(a-1)n}{l} (-\lambda)^{(a-1)n-l} H_{n-1}^{(an)}(x + l|\lambda).$$

Let us consider the following associated sequences:

$$\frac{1}{(1 - \lambda)^n} x H_{n-1}^{(n)}(x|\lambda) \sim (1, t(e^t - \lambda)), \quad p_n(x) \sim \left( 1, \left( \frac{1 - \lambda}{e^t - \lambda} \right)^a t \right), \quad (a \neq 0). \quad (2.7)$$

For  $x^n \sim (1, t)$ , by (1.11) and (2.7), we get

$$\begin{aligned} p_n(x) &= x \left( \frac{t}{t \left( \frac{1-\lambda}{e^t-\lambda} \right)^a} \right)^n x^{-1} x^n = x \left( \frac{e^t - \lambda}{1 - \lambda} \right)^{an} x^{n-1} \\ &= x \frac{1}{(1-\lambda)^{an}} \sum_{l=0}^{an} \binom{an}{l} (-\lambda)^{an-l} (x+l)^{n-1}. \end{aligned} \quad (2.8)$$

For  $n \geq 1$ , by (1.11) and (2.7), we get

$$\begin{aligned} p_n(x) &= x \left( \frac{t(e^t - \lambda)}{t \left( \frac{1-\lambda}{e^t-\lambda} \right)^a} \right)^n x^{-1} \frac{x}{(1-\lambda)^n} H_{n-1}^{(n)}(x|\lambda) \\ &= x \left( \frac{1}{1-\lambda} \right)^{(a+1)n} (e^t - \lambda)^{(a+1)n} H_{n-1}^{(n)}(x|\lambda). \end{aligned} \quad (2.9)$$

By (2.8) and (2.9), we get

$$\begin{aligned} &\sum_{l=0}^{an} \binom{an}{l} (-\lambda)^{an-l} (x+l)^{n-1} \\ &= \frac{1}{(1-\lambda)^n} (e^t - \lambda)^{(a+1)n} H_{n-1}^{(n)}(x|\lambda) \\ &= \frac{1}{(1-\lambda)^n} \sum_{l=0}^{(a+1)n} \binom{(a+1)n}{l} (-\lambda)^{(a+1)n-l} H_{n-1}^{(n)}(x+l|\lambda). \end{aligned} \quad (2.10)$$

Therefore, by (2.10), we obtain the following theorem.

**Theorem 2.2.** *For  $n \geq 1$  and  $a \in \mathbb{Z}_+ = \mathbb{N} \cup \{0\}$ , we have*

$$\sum_{l=0}^{an} \binom{an}{l} (-\lambda)^{-l} (x+l)^{n-1} = \frac{1}{(1-\lambda)^n} \sum_{l=0}^{(a+1)n} \binom{(a+1)n}{l} (-\lambda)^{n-l} H_{n-1}^{(n)}(x+l|\lambda).$$

Let us consider the following associated sequences:

$$(x)_n \sim (1, e^t - 1), \quad xH_{n-1}^{(an)}(x|\lambda) \sim \left( 1, t \left( \frac{e^t - \lambda}{1 - \lambda} \right)^a \right), \quad (a \neq 0). \quad (2.11)$$

By (1.11) and (2.11), we get

$$\begin{aligned} xH_{n-1}^{(an)}(x|\lambda) &= x \left( \frac{e^t - 1}{t \left( \frac{e^t - \lambda}{1 - \lambda} \right)^a} \right)^n x^{-1} (x)_n \\ &= x \left( \frac{e^t - 1}{t} \right)^n \left( \frac{1 - \lambda}{e^t - \lambda} \right)^{an} (x-1)_{n-1}. \end{aligned} \quad (2.12)$$

Replacing  $x$  by  $x + 1$ , we have

$$\begin{aligned}
 H_{n-1}^{(an)}(x+1|\lambda) &= \left(\frac{e^t-1}{t}\right)^n \left(\frac{1-\lambda}{e^t-\lambda}\right)^{an} \sum_{l=0}^{n-1} S_1(n-1, l) x^l \\
 &= \left(\frac{e^t-1}{t}\right)^n \sum_{l=0}^{n-1} S_1(n-1, l) H_l^{(an)}(x|\lambda) \\
 &= \sum_{l=0}^{n-1} \sum_{k=0}^l S_1(n-1, l) \frac{n!}{(k+n)!} S_2(k+n, n)(l)_k H_{l-k}^{(an)}(x|\lambda) \\
 &= \sum_{l=0}^{n-1} \sum_{k=0}^l S_1(n-1, l) S_2(k+n, n) \frac{\binom{l}{k}}{\binom{k+n}{n}} H_{l-k}^{(an)}(x|\lambda).
 \end{aligned} \tag{2.13}$$

Therefore, by (2.13), we obtain the following theorem.

**Theorem 2.3.** *For  $n \geq 1$ ,  $a \in \mathbb{Z}_+$ , we have*

$$H_{n-1}^{(an)}(x+1|\lambda) = \sum_{l=0}^{n-1} \sum_{k=0}^l S_1(n-1, l) S_2(k+n, n) \frac{\binom{l}{k}}{\binom{k+n}{n}} H_{l-k}^{(an)}(x|\lambda)$$

Let

$$\begin{aligned}
 x H_{n-1}^{(an)}(x|\lambda) &\sim \left(1, t \left(\frac{e^t-\lambda}{1-\lambda}\right)^a\right), \quad (a \neq 0), \\
 (x)_n &\sim (1, e^t - 1).
 \end{aligned} \tag{2.14}$$

Then, by (1.11) and (2.14), we get

$$\begin{aligned}
 (x)_n &= x \left(\frac{t \left(\frac{e^t-\lambda}{1-\lambda}\right)^a}{e^t-1}\right)^n x^{-1} x H_{n-1}^{(an)}(x|\lambda) \\
 &= x \left(\frac{t}{e^t-1}\right)^n \left(\frac{e^t-\lambda}{1-\lambda}\right)^{an} H_{n-1}^{(an)}(x|\lambda) \\
 &= x \left(\frac{t}{e^t-1}\right)^n x^{n-1} \\
 &= x B_{n-1}^{(n)}(x).
 \end{aligned} \tag{2.15}$$

and

$$(x)_n = \sum_{l=0}^n S_1(n, l) x^l = x \sum_{l=0}^{n-1} S_1(n, l+1) x^l, \quad (n \geq 1). \tag{2.16}$$

Therefore, by (2.15) and (2.16), we get

**Theorem 2.4.** *For  $n \geq 1$ ,  $0 \leq l \leq n-1$ , we have*

$$S_1(n, l+1) = \binom{n-1}{l} B_{n-1-l}^{(n)}.$$

From (2.15), we note that

$$\left(\frac{e^t-1}{t}\right)^n (x-1)_{n-1} = (1-\lambda)^{-an} (e^t-\lambda)^{an} H_{n-1}^{(an)}(x|\lambda), \quad (n \geq 1). \tag{2.17}$$

$$\begin{aligned}
LHS \text{ of (2.17)} &= \left( \frac{e^t - 1}{t} \right)^n \sum_{l=0}^{n-1} S_1(n-1, l) (x-1)^l \\
&= \sum_{l=0}^{n-1} S_1(n-1, l) \sum_{k=0}^l \frac{n!l!}{(k+n)!(l-k)!} S_2(k+n, n) (x-1)^{l-k} \\
&= \sum_{l=0}^{n-1} S_1(n-1, l) \sum_{k=0}^l \frac{\binom{l}{k}}{\binom{k+n}{n}} S_2(k+n, n) (x-1)^{l-k},
\end{aligned} \tag{2.18}$$

and

$$\begin{aligned}
RHS \text{ of (2.17)} &= (1-\lambda)^{-an} \sum_{l=0}^{an} \binom{an}{l} (-\lambda)^{an-l} e^{lt} H_{n-1}^{(an)}(x|\lambda) \\
&= (1-\lambda)^{-an} \sum_{l=0}^{an} \binom{an}{l} (-\lambda)^{an-l} H_{n-1}^{(an)}(x+l|\lambda).
\end{aligned} \tag{2.19}$$

Therefore, by (2.17), (2.18) and (2.19), we obtain the following theorem.

**Theorem 2.5.** *For  $n \geq 1$ ,  $a \in \mathbb{Z}_+$ , we have*

$$\begin{aligned}
&(1-\lambda)^{-an} \sum_{l=0}^{an} \binom{an}{l} (-\lambda)^{an-l} H_{n-1}^{(an)}(x+l|\lambda) \\
&= \sum_{l=0}^{n-1} \sum_{k=0}^l \frac{\binom{l}{k}}{\binom{k+n}{n}} S_1(n-1, l) S_2(k+n, n) (x-1)^{l-k}.
\end{aligned}$$

Let

$$p_n(x) \sim \left( 1, \left( \frac{1-\lambda}{e^t - \lambda} \right)^a t \right), \quad (x)_n \sim (1, e^t - 1), \quad (a \neq 0). \tag{2.20}$$

By (2.8), we have

$$\begin{aligned}
p_n(x) &= \left( \frac{1}{1-\lambda} \right)^{an} x \sum_{l=0}^{an} \binom{an}{l} (-\lambda)^{an-l} (x+l)^{n-1} \\
&= \left( \frac{1}{1-\lambda} \right)^{an} x \sum_{k=0}^{an} \binom{an}{k} (-\lambda)^{an-k} \sum_{l=0}^{n-1} \binom{n-1}{l} k^{n-1-l} x^l.
\end{aligned} \tag{2.21}$$

From (1.11) and (2.20), we have

$$\begin{aligned}
 (x)_n &= x \left( \frac{t \left( \frac{1-\lambda}{e^t - \lambda} \right)^a}{e^t - 1} \right)^n x^{-1} p_n(x) \\
 &= x \left( \frac{t}{e^t - 1} \right)^n \left( \frac{1-\lambda}{e^t - \lambda} \right)^{an} \left( \frac{1}{1-\lambda} \right)^{an} \sum_{k=0}^{an} \sum_{l=0}^{n-1} \binom{an}{k} \binom{n-1}{l} k^{n-1-l} (-\lambda)^{an-k} x^l \\
 &= \left( \frac{1}{1-\lambda} \right)^{an} x \left( \frac{t}{e^t - 1} \right)^n \sum_{k=0}^{an} \sum_{l=0}^{n-1} \binom{an}{k} \binom{n-1}{l} k^{n-1-l} (-\lambda)^{an-k} H_l^{(an)}(x|\lambda) \\
 &= \left( \frac{1}{1-\lambda} \right)^{an} x \sum_{k=0}^{an} \sum_{l=0}^{n-1} \sum_{m=0}^l \binom{an}{k} \binom{n-1}{l} \binom{l}{m} k^{n-1-l} (-\lambda)^{an-k} H_{l-m}^{(an)}(\lambda) B_m^{(n)}(x) \\
 &= \left( \frac{1}{1-\lambda} \right)^{an} x \sum_{k=0}^{an} \sum_{l=0}^{n-1} \sum_{m=0}^l \sum_{p=0}^m \binom{an}{k} \binom{n-1}{l} \binom{l}{m} \binom{m}{p} k^{n-1-l} (-\lambda)^{an-k} H_{l-m}^{(an)}(\lambda) B_{m-p}^{(n)} x^p \\
 &= \left( \frac{1}{1-\lambda} \right)^{an} x \sum_{p=0}^{n-1} \left\{ \sum_{k=0}^{an} \sum_{l=p}^{n-1} \sum_{m=p}^l \binom{an}{k} \binom{n-1}{l} \binom{l}{m} \binom{m}{p} k^{n-1-l} (-\lambda)^{an-k} H_{l-m}^{(an)}(\lambda) B_{m-p}^{(n)} \right\} x^p.
 \end{aligned} \tag{2.22}$$

Therefore, by (2.16) and (2.22), we obtain the following theorem.

**Theorem 2.6.** For  $n \geq 1$ ,  $a \in \mathbb{Z}_+$  and  $0 \leq p \leq n-1$ , we have

$$S_1(n, p+1) = \left( \frac{1}{1-\lambda} \right)^{an} \sum_{k=0}^{an} \sum_{l=p}^{n-1} \sum_{m=p}^l \binom{an}{k} \binom{n-1}{l} \binom{l}{m} \binom{m}{p} k^{n-1-l} (-\lambda)^{an-k} H_{l-m}^{(an)}(\lambda) B_{m-p}^{(n)}.$$

**Theorem 2.7.** For  $n \geq 0$ , we have

$$e^{-xt} (e^t - \lambda)^n = \sum_{k=0}^{\infty} \left( \sum_{l=0}^n \sum_{j=0}^k \binom{n}{l} (1-\lambda)^{n-l} \frac{\binom{k}{j}}{\binom{j+l}{l}} S_2(j+l, l) (-1)^{k-j} x^{k-j} \right) \frac{t^{k+l}}{k!}.$$

*Proof.* Note that

$$e^{-xt} (e^t - \lambda)^n = e^{-xt} (e^t - 1 + 1 - \lambda)^n = \sum_{l=0}^n \binom{n}{l} (1-\lambda)^{n-l} (e^t - 1)^l e^{-xt}, \tag{2.23}$$

and

$$\begin{aligned}
 (e^t - 1)^l e^{-xt} &= \sum_{k=0}^{\infty} \left( \sum_{j=0}^k \frac{l! k!}{(j+l)!(k-j)!} S_2(j+l, l) (-1)^{k-j} x^{k-j} \right) \frac{t^{k+l}}{k!} \\
 &= \sum_{k=0}^{\infty} \left( \sum_{j=0}^k \frac{\binom{k}{j}}{\binom{j+l}{l}} S_2(j+l, l) (-1)^{k-j} x^{k-j} \right) \frac{t^{k+l}}{k!}.
 \end{aligned} \tag{2.24}$$

From (2.23) and (2.24), we can derive Theorem 2.7.  $\square$

By (1.12) and Theorem 2.7, we get

$$\begin{aligned} A_n(x; b) &= x(x - bn)^{n-1} = x \left( \frac{1}{1-\lambda} \right)^{an} e^{-nbt} (e^t - \lambda)^{an} H_{n-1}^{(an)}(x|\lambda) \\ &= \frac{x}{(1-\lambda)^{an}} \sum_{l=0}^{an} \sum_{k=0}^{n-1-l} \sum_{j=0}^k \frac{\binom{an}{l} \binom{k}{j} \frac{(n-1)_{k+l}}{k!}}{\binom{j+l}{l}} (1-\lambda)^{an-l} S_2(j+l, l) (-1)^{k-j} (nb)^{k-j} H_{n-1-l-k}^{(an)}(x|\lambda). \end{aligned} \quad (2.25)$$

Therefore, by (2.25), we obtain the following theorem.

**Theorem 2.8.** *For  $n \geq 1$ ,  $a \in \mathbb{Z}_+$  and  $b \neq 0$ , we have*

$$(x - bn)^{n-1} = \sum_{l=0}^{an} \sum_{k=0}^{n-1-l} \sum_{j=0}^k \frac{\binom{an}{l} \binom{k}{j} \frac{(n-1)_{k+l}}{k!}}{\binom{j+l}{l} (1-\lambda)^l} S_2(j+l, l) (-1)^{k-j} (nb)^{k-j} H_{n-1-l-k}^{(an)}(x|\lambda).$$

Let us consider the Changhee polynomials of the second kind as follows:

$$\sum_{k=0}^{\infty} C_k(x|\lambda) \frac{t^k}{k!} = \frac{1}{1 + \lambda(1+t)} (1+t)^x. \quad (2.26)$$

From (1.10) and (2.26), we note that

$$C_k(x|\lambda) \sim (1 + \lambda e^t, e^t - 1). \quad (2.27)$$

Hence  $\lambda \in \mathbb{C}$  with  $\lambda \neq -1$ . Thus, by (2.27), we get

$$(1 + \lambda e^t) C_n(x|\lambda) = (x)_n \sim (1, e^t - 1), \quad (2.28)$$

and

$$(x)_n = x \left( \frac{t}{e^t - 1} \right)^n x^{-1} x^n = x \left( \frac{t}{e^t - 1} \right)^n x^{n-1} = x B_{n-1}^{(n)}(x). \quad (2.29)$$

Thus, by (2.28) and (2.29), we get

$$\begin{aligned} C_n(x|\lambda) &= \frac{1}{\lambda e^t + 1} x B_{n-1}^{(n)}(x) = \sum_{l=0}^n (-\lambda)^l e^{lt} \left( x B_{n-1}^{(n)}(x) \right) \\ &= \sum_{l=0}^n (-\lambda)^l (x+l) B_{n-1}^{(n)}(x+l). \end{aligned} \quad (2.30)$$

Let

$$t_n(x|\lambda) \sim \left( 1, \frac{t}{1 + \lambda(1+t)} \right). \quad (2.31)$$



Then, by (1.11) and (2.31), we get

$$\begin{aligned}
 t_n(x|\lambda) &= x \left( \frac{t}{\frac{t}{1+\lambda(1+t)}} \right)^n x^{-1} x^n = x(1+\lambda(1+t))^n x^{n-1} \\
 &= x \sum_{l=0}^n \binom{n}{l} \lambda^l (1+t)^l x^{n-1} = x \sum_{a=0}^n \binom{n}{a} \lambda^a \sum_{b=0}^{n-1} \binom{a}{b} t^b x^{n-1} \\
 &= \sum_{a=0}^n \sum_{b=0}^{n-1} \binom{n}{a} \binom{a}{b} \lambda^a (n-1)_b x^{n-b} = \sum_{a=0}^n \sum_{b=1}^n \lambda^a \binom{n}{a} \binom{a}{n-b} (n-1)_{n-b} x^b \\
 &= \sum_{a=0}^n \sum_{b=1}^n \lambda^a \binom{n}{a} \binom{a}{n-b} \frac{(n-1)!}{(b-1)!} x^b.
 \end{aligned} \tag{2.32}$$

Let us also consider the following associated sequence:

$$S_n(x|\mu) \sim \left( 1, \frac{t}{(1+t)^\mu} \right), \quad (\mu \in \mathbb{N}). \tag{2.33}$$

Then, by (1.11) and (2.33), we easily get

$$S_n(x|\mu) = \sum_{k=1}^n \binom{\mu n}{n-k} \frac{(n-1)!}{(k-1)!} x^k. \tag{2.34}$$

From (1.11), (2.32) and (2.33), we can derive

$$\begin{aligned}
 S_n(x|\mu) &= x \left( \frac{\frac{t}{1+\lambda(1+t)}}{\frac{t}{(1+t)^\mu}} \right)^n x^{-1} t_n(x|\lambda) = x \left( \frac{(1+t)^\mu}{1+\lambda(1+t)} \right)^n x^{-1} t_n(x|\lambda) \\
 &= x \left( \sum_{l=0}^{\infty} \frac{C_l(\mu|\lambda)}{l!} t^l \right)^n x^{-1} t_n(x|\lambda) \\
 &= x \sum_{l=0}^{\infty} \left\{ \sum_{l_1+\dots+l_n=l} \binom{l}{l_1, \dots, l_n} \left( \prod_{i=1}^n C_{l_i}(\mu|\lambda) \right) \right\} \frac{t^l}{l!} x^{-1} t_n(x|\lambda) \\
 &= x \sum_{l=0}^{\infty} \left\{ \sum_{l_1+\dots+l_n=l} \binom{l}{l_1, \dots, l_n} \left( \prod_{i=1}^n C_{l_i}(\mu|\lambda) \right) \right\} \frac{t^l}{l!} \left\{ \sum_{a=0}^n \sum_{b=1}^n \lambda^a \binom{n}{a} \binom{a}{n-b} \frac{(n-1)!}{(b-1)!} x^{b-1} \right\} \\
 &= x \sum_{a=0}^n \sum_{b=1}^n \sum_{l=0}^{b-1} \sum_{l_1+\dots+l_n=l} \binom{l}{l_1, \dots, l_n} \left( \prod_{i=1}^n C_{l_i}(\mu|\lambda) \right) \lambda^a \binom{n}{a} \binom{a}{n-b} \frac{(n-1)!}{(b-1)!} \frac{(b-1)_l}{l!} x^{b-1-l} \\
 &= \sum_{a=0}^n \sum_{b=1}^n \sum_{l=0}^{b-1} \sum_{l_1+\dots+l_n=l} \binom{l}{l_1, \dots, l_n} \left( \prod_{i=1}^n C_{l_i}(\mu|\lambda) \right) \lambda^a \binom{n}{a} \binom{a}{n-b} \frac{(n-1)!}{(b-1)!} \binom{b-1}{l} x^{b-l} \\
 &= \sum_{k=1}^n \left\{ \sum_{a=0}^n \sum_{b=k}^n \sum_{l_1+\dots+l_n=b-k} \binom{b-k}{l_1, \dots, l_n} \left( \prod_{i=1}^n C_{l_i}(\mu|\lambda) \right) \lambda^a \binom{n}{a} \binom{a}{n-b} \binom{b-1}{k-1} \frac{(n-1)!}{(b-1)!} \right\} x^k.
 \end{aligned} \tag{2.35}$$

Therefore, by (2.34) and (2.35), we obtain the following theorem.

**Theorem 2.9.** For  $n \geq 1$ ,  $1 \leq k \leq n$ ,  $b \neq 0$  and  $\mu, a \in \mathbb{Z}_+$ , we have

$$\frac{\binom{\mu n}{n-k}}{(k-1)!} = \sum_{a=0}^n \sum_{b=k}^n \sum_{l_1+\dots+l_n=b-k} \binom{n}{a} \binom{a}{n-b} \binom{b-k}{l_1, \dots, l_n} \binom{b-1}{k-1} \lambda^a \frac{1}{(b-1)!} \left( \prod_{i=1}^n C_{l_i}(\mu|\lambda) \right).$$

REMARK. From (1.1), we note that

$$\begin{aligned} \frac{1-\lambda}{e^t-\lambda} &= \frac{1-\lambda}{e^t-1+1-\lambda} = \frac{1}{1+\frac{e^t-1}{1-\lambda}} = \sum_{l=0}^{\infty} (-1)^l \left( \frac{e^t-1}{1-\lambda} \right)^l \\ &= \sum_{k=0}^{\infty} \left( \sum_{l=0}^k (-1)^l \left( \frac{1}{1-\lambda} \right)^l l! S_2(k, l) \right) \frac{t^k}{k!}, \end{aligned} \quad (2.36)$$

and

$$\frac{1-\lambda}{e^t-\lambda} = \sum_{n=0}^{\infty} H_n(\lambda) \frac{t^n}{n!}, \quad (2.37)$$

where  $H_n(\lambda)$  are the Frobenius-Euler numbers. By (2.36) and (2.37), we get

$$H_k(\lambda) = \sum_{l=0}^k \left( \frac{1}{\lambda-1} \right)^l l! S_2(k, l). \quad (2.38)$$

Let us consider the following associated sequences:

$$p_n(x) \sim \left( 1, \frac{1-\lambda}{e^t-\lambda} t \right), \quad x^n \sim (1, t). \quad (2.39)$$

Then, by (1.11) and (2.39), we get

$$p_n(x) = x \left( \frac{1}{1-\lambda} \right)^n (e^t - \lambda)^n x^{n-1} = \left( \frac{1}{1-\lambda} \right)^n x \sum_{k=0}^n \binom{n}{k} (-\lambda)^{n-k} (x+k)^{n-1}, \quad (2.40)$$

and

$$x^n = x \left( \frac{\left( \frac{1-\lambda}{e^t-\lambda} \right) t}{t} \right)^n x^{-1} p_n(x) = x \left( \frac{1-\lambda}{e^t-\lambda} \right)^n x^{-1} p_n(x). \quad (2.41)$$

Thus, by (2.40) and (2.41), we get

$$\begin{aligned} x^{n-1} &= \sum_{l=0}^{n-1} \left\{ \sum_{l_1+\dots+l_n=l} \binom{l}{l_1, \dots, l_n} H_{l_1}(\lambda) \cdots H_{l_n}(\lambda) \right\} \frac{t^l}{l!} \times \left\{ \sum_{k=0}^n \binom{n}{k} (-\lambda)^{n-k} \frac{(x+k)^{n-1}}{(1-\lambda)^n} \right\} \\ &= \frac{1}{(1-\lambda)^n} \sum_{k=0}^n \sum_{l=0}^{n-1} \sum_{l_1+\dots+l_n=l} \binom{l}{l_1, \dots, l_n} \left( \prod_{i=1}^n H_{l_i}(\lambda) \right) \binom{n}{k} \binom{n-1}{l} (-\lambda)^{n-k} (x+k)^{n-1-l}. \end{aligned}$$

## REFERENCES

- [1] S. Araci and M. Acikgoz, *A note on the Frobenius-Euler numbers and polynomials associated with Bernstein polynomials*, Adv. Stud. Contemp. Math., **22** (2012), 399-406.
- [2] S. Araci, M. Acikgoz and A. Esi, *A note on the  $q$ -Dedekind-type Daehee-Changhee sums with weight alpha arising from modified  $q$ -Genocchi polynomials with weight alpha*, arXiv:1211.2350.
- [3] S. Araci and M. Acikgoz, *Extended  $q$ -Dedekind-type Daehee-Changhee sums associated with Extended  $q$ -Euler polynomials*, arXiv:1211.1233.
- [4] S. Araci, E. Sen and M. Acikgoz, *A note on the modified  $q$ -Dedekind sums*, arXiv:1212.5837.

- [5] L. Carlitz, *Eulerian numbers and polynomials of higher order*, Duke Math. J., **27** (1960), 401-423.
- [6] I. N. Cangul, V. Kurt, H. Ozden and Y. Simsek, *On the higher-order  $w$ - $q$ -Genocchi numbers*, Adv. Stud. Contemp. Math. **19** (2009), no. 1, 39-57.
- [7] R. Dere and Y. Simsek, *Applications of umbral algebra to some special polynomials*, Adv. Stud. Contemp. Math. **22** (2012), no. 3, 433-438.
- [8] Q. Fang and T. Wang, *Umbral calculus and invariant sequences*, Ars Combin., **101** (2011) 257-264.
- [9] D. S. Kim, T. Kim, S. H. Lee and S. H. Rim, *Frobenius-Euler polynomials and umbral calculus in the  $p$ -adic case*, Adv. Difference Equ. 2012, 2012:222.
- [10] D. S. Kim and T. Kim, *Some new identities of Frobenius-Euler numbers and polynomials*, J. of Inequ. and Appl., 2012, 2012:307.
- [11] D. S. Kim and T. Kim, *Applications of Umbral Calculus Associated with  $p$ -Adic Invariant Integrals on  $\mathbb{Z}_p$* , Abstract and Applied Analysis 2012 (2012), Article ID 865721, 12 pages.
- [12] D. S. Kim and T. Kim, *Some identities of Frobenius-Euler polynomials arising from umbral calculus*, Adv. Difference Equ. 2012, 2012:196.
- [13] T. Kim, *Identities involving Frobenius-Euler polynomials arising from non-linear differential equations*, J. Number Theory, **132** (2012), no. 1, 2854-2865.
- [14] T. Kim, *Some identities on the  $q$ -Euler polynomials of higher order and  $q$ -Stirling numbers by the fermionic  $p$ -adic integral on  $\mathbb{Z}_p$* , Russ. J. Math. Phys., **16** (2009), no. 4, 484-491.
- [15] T. Kim, *An identity of the symmetry for the Frobenius-Euler polynomials associated with the fermionic  $p$ -adic invariant  $q$ -integrals on  $\mathbb{Z}_p$* , Rocky Mountain J. Math., **41** (2011), no. 1, 239-247.
- [16] T. J. Robinson, *Formal calculus and umbral calculus*, Electron. J. Combin., **17** (2010), no. 1, Research Paper 95, 31 pp.
- [17] S. Roman, *More on the umbral calculus, with emphasis on the  $q$ -umbral calculus*, J. Math. Anal. Appl., **107** (1985), 222-254.
- [18] S. Roman, *The umbral calculus*, Dover Publ. Inc. New York, 2005.
- [19] C. Ryoo, *A note on the Frobenius-Euler polynomials*, Proc. Jangjeon Math. Soc., **14** (2011), no. 4, 495-501.
- [20] C. S. Ryoo and R. P. Agarwal, *Calculating zeros of the Frobenius-Euler polynomials*, Neural Parallel Sci. Comput., **17** (2009), no. 3, 351-361.
- [21] C. S. Ryoo and R. P. Agarwal, *Exploring the multiple Changhee  $q$ -Bernoulli polynomials*, Int. J. Comput. Math., **82** (2005), no. 4, 483-493.
- [22] Y. Simsek, O. Yurekli and V. Kurt, *On interpolation functions of the twisted generalized Frobenius-Euler numbers*, Adv. Stud. Contemp. Math., **15** (2007), no. 2, 187-194.
- [23] K. Shiratani, *On Euler numbers*, Mem. Fac. Sci. Kyushu Univ. Ser. A, **27** (1973), 1-5.

<sup>1</sup> DEPARTMENT OF MATHEMATICS, KWANGWOON UNIVERSITY, SEOUL 139-701, REPUBLIC OF KOREA.

*E-mail address:* `tkkim@kw.ac.kr`

<sup>2</sup> DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 121-742, REPUBLIC OF KOREA.

*E-mail address:* `dskim@sogang.ac.kr`